

NONNEGATIVE POLYNOMIALS FROM VECTOR BUNDLES ON REAL CURVES

ROGER BIELAWSKI

ABSTRACT. We observe that the E -resultant of a very ample rank 2 vector bundle E on a real projective curve (with no real points) is nonnegative when restricted to the space of real sections. Moreover, we show that if E has a section vanishing at exactly two points and the degree d of E satisfies $d(d-6) \geq 4(g-1)$, then this polynomial cannot be written as a sum of squares.

1. INTRODUCTION

Ever since Hilbert's seminal 1888 paper [5] it has been known that there exist real polynomials in n variables ($n \geq 3$), which are nonnegative on all of $\mathbb{R}^n$, but cannot be written as a sum of squares of polynomials. Hilbert's proof was nonconstructive and the first explicit example was found only in 1967. The subject of nonnegative polynomials has become a rapidly evolving area of real algebraic geometry, with links and applications to semidefinite programming and polynomial optimization problems: see, e.g., the monograph [6].

In this note, we are going to show that a class of nonnegative polynomials, which cannot be written as sums of squares, corresponds to very ample rank 2 vector bundles on real projective curves with no real points (the vector bundles must satisfy some additional constraints; see below). The simplest vector bundle satisfying the necessary conditions is $\mathcal{O}(3) \oplus \mathcal{O}(3)$ on $\mathbb{CP}^1$ equipped with the antipodal map, and we compute explicitly the corresponding nonnegative polynomial which is not a sum of squares (Example 3.3). It is of degree 6 in 8 variables, with 224 terms.

Let us explain the correspondence between vector bundles and polynomials in greater detail. Let E be a globally generated nontrivial vector bundle of rank 2 on a smooth connected projective curve C (over $\mathbb{C}$). The cone

$$V(E) = \{s \in H^0(C, E); \exists_{z \in C} s(z) = 0\}$$

of sections of E which vanish somewhere is an irreducible hypersurface of $H^0(C, E)$, and hence the zero set of an irreducible homogeneous polynomial R_E . Following Gelfand, Kapranov, and Zelevinsky [4, Ch. 3], we call R_E the E -resultant.

Suppose now, in addition, that C is a real curve, i.e. C is equipped with an antiholomorphic involution σ , and that E is a σ -bundle, i.e. there is an antilinear bundle isomorphism $\tau : E \rightarrow E$ which covers σ and satisfies $\tau^2 = 1$. We can then restrict R_E to the real vector space $H^0(C, E)^\tau$ of τ -invariant sections and obtain (after multiplying by a constant) a real polynomial ρ_E on $H^0(C, E)^\tau$.

The real polynomial ρ_E is the subject of this note. It is of course determined only up to a nonzero multiple, and we can assume that ρ_E takes at least one positive value. Our main result is:

Theorem 1. *Let C be a smooth connected projective curve of genus g , equipped with an antiholomorphic involution σ with no fixed points. Let E be a rank 2 and degree d very ample σ -bundle on C . Then:*

- (i) ρ_E is a nonnegative homogeneous polynomial of degree d ;
- (ii) ρ_E vanishes on a set of codimension 2, unless E is $\mathcal{O}(1) \oplus \mathcal{O}(1)$ on $\mathbb{P}^1$. At a generic point of $\rho_E^{-1}(0)$, the rank of the Hessian of ρ_E is 4.
- (iii) If $d(d-6) \geq 4(g-1)$ and E has a section which vanishes at exactly two points, then ρ_E cannot be written as a sum of squares of polynomials.

Remark 1.1. A large class of curves equipped with an antiholomorphic involution with no fixed points is obtained by considering curves in noncompact complex surfaces equipped with such an involution, e.g. the total space of $\mathcal{O}(2k)$ on $\mathbb{P}^1$, or $\mathbb{P}^1 \times \overline{\mathbb{P}^1}$ minus the diagonal.

Remark 1.2. If E is nonspecial, i.e. $h^1(E) = 0$, then the numerical condition $d(d-6) \geq 4(g-1)$ is automatically satisfied if $g \geq 1$ (and if $g = 0$, $d \geq 6$). This follows by writing $d = 2(g-1) + n$, $n = h^0(E)$, and observing that $d(d-6) \geq 4(g-1)$ if $n \geq 6$. The case $n < 6$ cannot occur. Indeed, n must be even (as d is even) and there are no very ample rank 2 vector bundles with $h^0(E) = 2, 4$ on curves of positive genus (see, e.g., the proof of Lemma 2.1 below).

Remark 1.3. The condition that E has a section vanishing at exactly two points is satisfied, for example, if E is 2-very ample, i.e. the evaluation map $H^0(C, E) \rightarrow H^0(C, E \otimes \mathcal{O}_D)$ is surjective for any effective divisor D of length 3.

2. PROOF OF THEOREM 1

We shall first prove statements (i) and (ii). The fact that $\deg \rho_E = \deg E$ follows from Theorem 3.10 in Chapter 3 of [4]. The following lemma proves the first claim in statement (ii).

Lemma 2.1. *Let E be a rank 2 very ample σ -bundle on a real projective curve C with no real points. Then the codimension of $V(E)^\tau$ is two, unless E is $\mathcal{O}(1) \oplus \mathcal{O}(1)$ on $\mathbb{P}^1$.*

Proof. Suppose first that $h^0(E) = 4$. Then $S = \mathbb{P}V(E)$ is a geometrically ruled (by the $S_x = \mathbb{P}\{s \in H^0(E); s(x) = 0\}$) surface in $\mathbb{P}^3$. It follows that $b_2(S) = 2$ [2, Prop. III.21], and consequently, the degree of R_E is 2. Thus S is a quadric and $E \simeq \mathcal{O}(1) \oplus \mathcal{O}(1)$ on $\mathbb{P}^1$.

We can therefore assume that $n = h^0(E) > 4$. E being very ample is equivalent to $\dim H^0(C, E(-x-y)) = n-4$ for any $x, y \in C$. A τ -invariant section, which vanishes at an $x \in C$, must also vanish at $\sigma(x)$. Therefore, for any $x \in C$,

$$V(E)_x^\tau = \{s \in H^0(E)^\tau; s(x) = 0\} = \{s \in H^0(E); s(x) = s(\sigma(x)) = 0\}^\tau,$$

has real codimension 4 in $H^0(E)^\tau$. Let us write C/σ for the unoriented real surface obtained as the quotient of C by σ . Points of C/σ can be identified with divisors $x + \sigma(x)$ on C . We consider the incidence variety

$$Y = \{(x + \sigma(x), s) \in C/\sigma \times V(E)^\tau; s(x) = s(\sigma(x)) = 0\},$$

and the two projections p_1, p_2 onto C/σ and onto $V(E)^\tau$, respectively. The fibres of p_1 are the $V(E)_x^\tau$ and have real dimension $n-4$, which implies that Y has dimension $n-2$. The generic fibre of p_2 is finite and so $\dim_{\mathbb{R}} V(E)^\tau = \dim_{\mathbb{R}} Y = n-2$. $\square$

The following simple lemma implies that ρ_E must be nonnegative:

Lemma 2.2. *Let f be a real polynomial in n variables. If f changes sign, then $f^{-1}(0)$ has codimension 1 somewhere.*

Proof. After a linear change of variables, we can assume that there are $a \in \mathbb{R}^{n-1}$, $y_1, y_2 \in \mathbb{R}$, such that $f(a, y_1) < 0 < f(a, y_2)$. Therefore, for any x in a small neighbourhood U_a of a , we have $f(x, y_1) < 0 < f(x, y_2)$, and the mean value theorem applied to $F_x(t) = f(x, t)$ shows that, for any $x \in U_a$, there exists a $y_x \in \mathbb{R}$ such that $f(x, y_x) = 0$. $\square$

The statement about the Hessian of ρ_E follows from the following two facts:

1. The projective dual of $\mathbb{P}V(E)$ is $\mathbb{P}(E^*)$ embedded in $\mathbb{P}(W^*)$, $W = H^0(C, E)$, via the projective embedding defined by the line bundle $\mathcal{O}(1)$ on $\mathbb{P}(E^*)$. This is shown in [4], Theorem 3.11 in Chapter 3.
2. Segre's dimension formula [9] (see also [4, Ch. 1, Thm. 5.3]).

2.1. The variety $V^{[2]}(E)$. In order to prove statement (iii) of Theorem 1, we shall consider the variety $V^{[2]}(E) \subset V(E)$ of sections which have at least 2 zeros (counting multiplicities).

Lemma 2.3. *Let E be a very ample vector bundle of rank 2 on a smooth connected projective curve C . The variety $\mathbb{P}V^{[2]}(E)$ is irreducible and nonempty unless E is $\mathcal{O}(1) \oplus \mathcal{O}(1)$ on $\mathbb{P}^1$. Moreover, $\mathbb{P}V^{[2]}(E)$ is the singular locus of $\mathbb{P}V(E)$.*

Proof. As in the proof of Lemma 2.1, we can assume that $h^0(E) > 4$. This time we consider the incidence variety

$$Y = \{(D, s) \in C_2 \times V^{[2]}(E); s(D) = 0\},$$

where $C_2 = C^2/\mathbb{Z}_2$. Let $p_1 : Y \rightarrow C_2$ and $p_2 : Y \rightarrow V^{[2]}(E)$ be the two projections. Y is the total space of a vector bundle over an irreducible variety, hence irreducible. Therefore $V^{[2]}(E)$ is irreducible and of codimension 2 (cf. the proof of Lemma 2.1).

To show that $\mathbb{P}V^{[2]}(E)$ is the singular locus of $\mathbb{P}V(E)$, consider the incidence variety

$$\tilde{Y} = \{(x, [s]) \in C \times \mathbb{P}V(E); s(x) = 0\}.$$

The projection p_2 onto $\mathbb{P}V(E)$ is ramified over $\mathbb{P}V^{[2]}(E)$, which implies the inclusion $\mathbb{P}V(E)_{\text{sing}} \subset \mathbb{P}V^{[2]}(E)$. On the other hand, as mentioned at the end of the last subsection, $\mathbb{P}V(E)$ is the projective dual of $X = \mathbb{P}(E^*)$ embedded in $\mathbb{P}(W^*)$, $W = H^0(C, E)$, via the projective embedding defined by the line bundle $\mathcal{O}(1)$ on $\mathbb{P}(E^*)$. Thus, the hyperplane H corresponding to $s \in H^0(C, E) \simeq H^0(X, \mathcal{O}(1))$ is tangent to X at two points (or has a tangency point of order 2) if and only if $s \in V^{[2]}(E)$. Therefore $\mathbb{P}V^{[2]}(E) \subset \mathbb{P}V(E)_{\text{sing}}$. $\square$

Lemma 2.4. *Let E be a very ample rank 2 σ -bundle on a real smooth connected projective curve C with no real points. If ρ_E is a sum of squares of polynomials, then $\deg V^{[2]}(E) < (\deg E)^2/4$.*

Proof. The inequality is clearly true for $E = \mathcal{O}(1) \oplus \mathcal{O}(1)$ on $\mathbb{P}^1$, so we exclude this case. Suppose that $\rho_E = \sum_{i=1}^m p_i^2$ for real polynomials $p_i(x)$ (p_i is not a constant multiple of p_j for $i \neq j$). Then the E -resultant is of the same form: $R_E(z) = \sum_{i=1}^m p_i^2(z)$. From the previous lemma we can conclude that the subset P of $H^0(C, E)$ defined by $p_i(z) = 0$, $i = 1, \dots, m$, is contained in $V^{[2]}(E)$. On the

other hand, the variety $V(E)^\tau$ is given by the equations $p_i(x) = 0$, $i = 1, \dots, m$ and it has codimension 2, owing to Lemma 2.1. Therefore P has codimension at most 2 and, as $V^{[2]}(E)$ is irreducible, $P = V^{[2]}(E)$. The degree of each p_i is at most $d/2$, and since $V^{[2]}(E)$ has codimension 2, it follows that its degree is at most $(d/2)^2$. Moreover, the irreducibility of $V(E)$ implies that $m \geq 3$, so the inequality is strict. $\square$

Theorem 1(iii) will now follow from:

Proposition 2.5. *Let C be a curve of genus g and E a rank 2 very ample vector bundle on C . Let $d = \deg E$ and suppose that E has a section with exactly 2 zeros. Then the degree of $V^{[2]}(E)$ is $\frac{d(d-3)}{2} + 1 - g$.*

In order to prove this, we need to discuss certain natural vector bundles on symmetric powers of C .

2.2. Tautological vector bundles on symmetric products and $\deg V^{[2]}(E)$. Let E be a vector bundle of rank r on an algebraic curve C . For any $k \geq 1$ one obtains a vector bundle $E^{[k]}$ of rank kr on the k -th symmetric product $C_k = C^k/\Sigma_k$. Consider

$$C \xleftarrow{q} C \times C_{k-1} \xrightarrow{p} C_k,$$

where q is the projection and p is the symmetrisation map. Then $E^{[k]} = p_* q^* E$. This construction goes back to Schwarzenberger [8]. Mattuck [7] has shown that $E^{[k]}$ can be also constructed as the Σ_k -invariant part of $f_*(q_1^* E \oplus \dots \oplus q_k^* E)$, where $q_i : C^k \rightarrow C$ is the projection onto the i -th factor and $f : C^k \rightarrow C_k$ is the symmetrisation. The bundle $E^{[k]}$ was called *symmetrization of E* by Mattuck, but the term “*tautological bundle*” seems to have been accepted in more recent literature.

The fibre of $E^{[k]}$ at $D \in C_k$ is $H^0(C, E \otimes \mathcal{O}_D)$. The Künneth formula implies easily that $H^0(C_k, E^{[k]}) \simeq H^0(C, E)$, and so we have the natural evaluation map

$$(2.1) \quad H^0(C, E) \rightarrow E^{[k]},$$

which is surjective at $D \in C_k$ if and only if the evaluation morphism

$$(2.2) \quad H^0(C, E) \rightarrow H^0(C, E \otimes \mathcal{O}_D)$$

is surjective. In particular, if E is very ample, then $E^{[2]}$ is globally generated. Moreover, the isomorphism $H^0(C_2, E^{[2]}) \simeq H^0(C, E)$ induces an isomorphism $V^{[2]}(E) \simeq V(E^{[2]})$ with the variety of sections of $E^{[2]}$ vanishing somewhere. The assumption on E having a section vanishing at exactly two points is equivalent to generic section in $V(E^{[2]})$ having exactly one zero. In this case, the results of Geertsens [3, Formula 11] imply that the degree of $V^{[2]}(E) \simeq V(E^{[2]})$ is $c_2(E^{[2]})[C_2]$.

The Chern character of the tautological bundle $E^{[k]}$ has been computed by Mattuck [7]¹, and we simply apply his results (see also [1, §VIII.2]). Let $\delta \in H^2(C_2, \mathbb{Q})$ be the Poincaré dual of the image of the diagonal $\{(p, p)\}$ in C_2 (i.e. of $\{2p\} \subset C_2$). Let $x \in H^2(C_2, \mathbb{Q})$ be the Poincaré dual of $p + C$, and 1 the Poincaré dual of $[C_2]$. These satisfy the following relations:

$$x^2 = 1, \quad x\delta = 2, \quad \delta^2 = 4(1 - g).$$

¹Mattuck only considers line bundles E , but, owing to a standard argument, this is sufficient to compute the Chern character for higher rank bundles.

The Chern character of $E^{[2]}$, where $\text{rank } E = r$ and $\deg(E) = d$, is

$$\text{ch}(E^{[2]}) = d(1 - e^{-x}) - r(g - 1) + r \left((g + 1)(1 + x) - \frac{\delta}{2} \right) e^{-x}.$$

Hence:

$$(2.3) \quad c_1(E^{[2]}) = dx - r \frac{\delta}{2}, \quad c_2(E^{[2]}) = \frac{d(d + 1 - 2r)}{2} - \frac{r(r - 1)}{2}(g - 1).$$

Thus, for a rank 2 bundle E of degree d , the degree of the variety $V^{[2]}(E)$ is $\frac{d(d-3)}{2} + 1 - g$, which proves Proposition 2.5 and, consequently, Theorem 1.

3. EXAMPLES

We discuss the case of $C = \mathbb{P}^1$, equipped with the antipodal map σ . If z is the affine coordinate on $\mathbb{P}^1$, then $\sigma(z) = -1/\bar{z}$. The involution σ induces an anti-linear map τ on $\mathcal{O}(k)$, which squares to 1 if k is even, and to -1 if k is odd. For $k \geq 0$, its effect on sections, i.e. on polynomials of degree k , is:

$$(3.1) \quad \tau \left(\sum_{i=0}^k c_i z^i \right) = \sum_{i=0}^k (-1)^{i+1} \overline{c_{k-i}} z^i.$$

It follows that the rank 2 σ -bundles on $\mathbb{P}^1$ are $\mathcal{O}(2k) \oplus \mathcal{O}(2l)$ with diagonal action of τ , or $\mathcal{O}(2k+1) \oplus \mathcal{O}(2k+1)$ with $\tau(u, v) = (\tau(v), -\tau(u))$. The polynomial R_E is the classical resultant $R(f, g)$ of two polynomials. We compute the real polynomial ρ_E for $E = \mathcal{O}(i) \oplus \mathcal{O}(i)$, $i = 1, 2, 3$:

3.1. $E = \mathcal{O}(1) \oplus \mathcal{O}(1)$. In this case, a real section of E is a pair $f = a_0 + a_1 z$, $g = \overline{a_1} - \overline{a_0} z$. The resultant $R(f, g)$ is $|a_1|^2 + |a_0|^2$, which is the quadratic polynomial $x_1^2 + x_2^2 + x_3^2 + x_4^2$ on $\mathbb{R}^4$.

3.2. $E = \mathcal{O}(2) \oplus \mathcal{O}(2)$. Real sections of E are pairs $f = a + rz - \bar{a}z^2$, $g = b + sz - \bar{b}z^2$, $a, b \in \mathbb{C}$, $r, s \in \mathbb{R}$, and the classical resultant of two such polynomials is

$$R(f, g) = |as - br|^2 - (a\bar{b} - \bar{a}b)^2.$$

After a linear change of coordinates, we can write this as the polynomial

$$\rho_E(x_1, x_2, x_3, y_1, y_2, y_3) = (x_1 y_2 - x_2 y_1)^2 + (x_1 y_3 - x_3 y_1)^2 + (x_2 y_3 - x_3 y_2)^2$$

on $\mathbb{R}^6$.

3.3. $E = \mathcal{O}(3) \oplus \mathcal{O}(3)$. This is the first case where Theorem 1 implies that ρ_E is not a sum of squares of polynomials.

Sections of E are pairs of polynomials $f = \sum_{i=0}^3 a_i z^i$, $g = \sum_{i=0}^3 b_i z^i$ of degree 3 and $R_E = R(f, g)$ can be computed from the Cayley-Bezout formula [4, §12.1]:

$$(3.2) \quad R(f, g) = \det \begin{pmatrix} [3, 0] & [3, 1] & [3, 2] \\ [2, 0] & [3, 0] + [2, 1] & [3, 1] \\ [1, 0] & [2, 0] & [3, 0] \end{pmatrix},$$

where $[i, j] = a_i b_j - b_i a_j$. We can identify $H^0(C, E)^\tau$ with pairs

$$f = a_0 + a_1 z + a_2 z^2 + a_3 z^3, \quad g = \overline{a_3} - \overline{a_2} z + \overline{a_1} z^2 - \overline{a_0} z^3.$$

It follows that on $H^0(C, E)^\tau$ we have

$$[3, 0] = |a_3|^2 + |a_0|^2, \quad [2, 1] = -|a_2|^2 - |a_1|^2,$$

$$[3, 2] = \overline{[1, 0]} = a_3\overline{a_1} + a_2\overline{a_0}, \quad [3, 1] = -\overline{[2, 0]} = -a_3\overline{a_2} + a_1\overline{a_0}.$$

Writing

$$r = [3, 0], \quad s = -[2, 1], \quad u = [3, 1], \quad v = [3, 2],$$

we obtain

$$(3.3) \quad \rho_E(f) = R(f, \sigma(f)) = r^2(r - s) + 2r|u|^2 - (r - s)|v|^2 + u^2\bar{v} + \bar{u}^2v.$$

It is a polynomial of degree 6 in 8 real variables, with 224 terms. Using the identity $|u|^2 + |v|^2 = rs$, we can eliminate s (if $r \neq 0$), and rewrite ρ_E in the following (nonpolynomial) form, which makes its nonnegativity obvious:

$$(3.4) \quad \rho_E(f) = \frac{1}{r} \left((r^2 - |v|^2)^2 + |r\bar{u} + u\bar{v}|^2 \right).$$

This also shows that ρ_E vanishes if and only if $r\bar{u} + u\bar{v} = 0$, i.e. if

$$(|a_0|^2 + |a_3|^2)(a_0\overline{a_1} - a_2\overline{a_3}) = (a_3\overline{a_2} - a_1\overline{a_0})(a_0\overline{a_2} + a_1\overline{a_3}).$$

REFERENCES

- [1] E. Arbarello, M. Cornalba, P.A. Griffiths, and J. Harris, *Geometry of algebraic curves. Vol. I*, Springer, New York, 1985.
- [2] A. Beauville, *Complex algebraic surfaces*, CUP, Cambridge, 1996.
- [3] J.A. Geerts, ‘Degeneracy loci of vector bundle maps and ampleness’, *Math. Scand.* 90 (2002), 13–34.
- [4] I.M. Gelfand, M.M. Kapranov, and A.V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Birkhäuser, Boston, 1994.
- [5] D. Hilbert, ‘Über die Darstellung definiter Formen als Summe von Formenquadraten’, *Math. Ann.* 32 (1888), 342–350.
- [6] M. Marshall, *Positive polynomials and sums of squares*, AMS, Providence, RI, 2008.
- [7] A. Mattuck, ‘Secant bundles on symmetric products’, *Amer. J. Math.* 87 (1965), 779–797.
- [8] R.L.E. Schwarzenberger, ‘The secant bundle of a projective variety’, *Proc. London Math. Soc.* (3), 14 (1964), 369–384.
- [9] B. Segre, ‘Bertini forms and Hessian matrices’, *J. London Math. Soc.* 26 (1951), 164–176.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UK